



TECHICAL NOTE

PLANT MATERIALS

23

FORAGE QUALITY ANALYSIS OF SEVERAL PULLMAN PLANT MATERIALS CENTER RELEASES.

by

Mark Stannard and Clarence Kelley
Spokane, Washington

October 1993

Forage quantity and quality are two very important factors which concern livestock producers. Forage quantity is commonly estimated by producers every year and livestock feed requirements are based on these estimates. Forage quality on the other hand is not as easily determined. "Ballpark estimates" are commonly used to estimate forage quality of pasture and range plants. Most producers would agree that "ballpark estimates" are inadequate and better estimates are needed.

Several plant releases made by the Pullman PMC were analyzed for feed quality by the *International Feedstuffs Institute* as part of a nation-wide project. Forage samples were taken at various stages of plant development from plants growing at the Pullman PMC in 1973, 1976, and 1980. Figure 1 and Table 1 show the various components of forage dry matter. Over fifteen separate feed analyses were completed and a printout was provided to the center. An abridged report of those results is shown in Table 2.

Forage quality varied considerably between growth stage and species. Quality of the grasses tested is best in the vegetative stages and decreases considerably with reproductive growth. Regrowth of pasture grasses such as 'Latar' orchardgrass and 'Alta' tall fescue exhibited higher quality than samples tested at reproductive stages.

Crude protein varied considerably with the stage of growth but varied less so between species sampled at the same stage of growth. For example, crude protein for 'Secar' bluebunch wheatgrass varied from 23.4 to 7.0% for the early vegetative stage and dough stage, respectively. Crude protein measured for samples collected in the late bloom stage ranged from 6.4 to 10.9% for 'Alkar' tall wheatgrass and 'Latar' orchardgrass, respectively. Crude protein may also vary between cultivars of a given species. However, the value of breeding cultivars for protein content is of doubtful value since protein content can be manipulated by stage of growth at harvest, (Howarth and Goplen, 1983).

The United States Department of Agriculture (USDA) prohibits discrimination in all its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write the USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C., 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

'Cascade' birdsfoot trefoil exhibited excellent forage quality at the mid bloom stage for most parameters with the exception of the lignin content, 12.4%. Alfalfa sampled at the same growth stage in comparison showed an average lignin content of 5.5% in the 1971 Atlas of Nutritional Data on United States and Canadian Feeds. Lignin is an important indigestible component of forage and is negatively associated with forage quality. Lignin content generally increases with plant maturation.

Table 1. Classification of forage fractions using the Van Soest method.

| Fraction | Components | -----Nutritional Availability----- | |
|-----------------|------------------------|------------------------------------|--------------|
| | | Ruminant | Non-ruminant |
| Cell Contents | Sugars, Starch, pectin | Complete | Complete |
| | Soluble carbohydrates | Complete | Complete |
| | Protein. Non-protein N | High | High |
| | Lipids (fats) | High | High |
| | Other solubles | High | High |
| Cell Wall (NDF) | Hemicellulose | Partial | Low |
| | Cellulose | Partial | Low |
| | Heat Damaged Protein | Indigestible | Indigestible |
| | Lignin | Indigestible | Indigestible |
| | Silica | Indigestible | Indigestible |

Source: Van Soest (1967)

In a similar study, the Pullman PMC provided forage samples of several accessions of Siberian wheatgrass and 'Nordan' crested wheatgrass to the WSU Forage Lab for analysis in 1992. Samples were collected at Central Ferry, Washington and taken at: 1) early vegetative growth, 2) late vegetative growth to boot, and 3) seed filling stages. Crude protein, Acid Detergent Fiber (ADF) and Acid Detergent Lignin (ADL) of 'Nordan' and 'P-27' Siberian wheatgrass are shown in Table 3. Clearly, forage quality diminishes as the two cultivars mature.

While the results of the analyses are good estimates, it is important to realize that these analyses results are for specific cultivars which were sampled at the PMC and Central Ferry at specific dates. Climatic conditions, soil conditions, and cultivar type all effect feed quality and quantity.

References


The Atlas of Nutritional Data on United States and Canadian Feeds (1971) published by the National Academy of Sciences.

Howarth, R.E. and B.P. Goplen. 1983. Improvement of forage quality through production management and plant breeding. Can. J. Plant Sci. 63:895-902.

Pioneer Forage Manual - A Nutritional Guide. 1990. Pioneer Hi-Bred International, Inc.

Van Soest, P.J. 1967. Development of a comprehensive system of feed analyses and its application to forages. J. Animal Sci. 26:119-128.

Figure 1. The detergent (Van Soest) procedure to partition forages.



Source: Pioneer Forage Manual - A Nutritional Guide. 1990.
Pioneer Hi-Bred International, Inc.

Table 2. SHORT SYNOPSIS OF THE FORAGE QUALITY DATA PRODUCED BY DR. FONNESBECK OF THE INTERNATIONAL FEEDSTUFFS INSTITUTE.

| CULTIVAR | DRY | | CRUDE | | HEMICELL- ⁴ | ACID | LIGNIN ⁶ | | Ca | Mg | P |
|-----------------------------|----------------|------------------|----------------------|------------------|------------------------|---------------------|---------------------|------------------------|------|-----|-----|
| STAGE OF GROWTH | MATTER | ASH ¹ | PROTEIN ² | NDF ³ | ULOSE | INSOL. ⁵ | | CELLULOSE ⁷ | | | |
| DATE SAMPLED | ------(%)----- | | | | | | | | | | |
| LATAR | | | | | | | | | | | |
| Orchardgrass | | | | | | | | | | | |
| late veg | --- | --- | 16.9 | 58.1 | 19.2 | 1.7 | 5.1 | 29.4 | -- | -- | -- |
| 5-10-76 | | | | | | | | | | | |
| mid bloom | --- | 6.9 | 7.0 | --- | --- | --- | --- | --- | .98 | .33 | .12 |
| 6-14-73 | | | | | | | | | | | |
| late bloom | 33.3 | 9.9 | 10.9 | 70.7 | 20.5 | 3.2 | 7.6 | 36.4 | .60 | .28 | .12 |
| 7-7-80 | | | | | | | | | | | |
| regrowth | --- | --- | 13.1 | 60.1 | 14.7 | 1.5 | 8.7 | 32.4 | -- | -- | -- |
| 8-2-73 | | | | | | | | | | | |
| SECAR | | | | | | | | | | | |
| Bluebunch Wheatgrass | | | | | | | | | | | |
| early veg | 36.0 | 16.4 | 23.4 | 53.9 | 15.6 | 6.9 | 3.4 | 25.3 | 1.04 | .42 | .29 |
| 3-10-80 | | | | | | | | | | | |
| late veg | 30.5 | 9.2 | 14.7 | 72.1 | 21.2 | 3.3 | 7.0 | 37.7 | .56 | .19 | .22 |
| 5-26-80 | | | | | | | | | | | |
| mid bloom | 49.8 | 8.0 | 9.1 | 71.5 | 15.3 | 3.5 | 9.6 | 40.2 | .47 | .13 | .16 |
| 7-7-80 | | | | | | | | | | | |
| late bloom | 41.8 | 9.0 | 10.5 | 73.5 | 19.0 | 4.9 | 8.9 | 37.8 | .51 | .10 | .18 |
| 7-10-80 | | | | | | | | | | | |
| dough | 54.2 | 7.0 | 7.0 | 71.7 | 17.8 | 4.0 | 10.1 | 36.8 | .39 | .10 | .14 |
| 7-18-80 | | | | | | | | | | | |
| WHITMAR | | | | | | | | | | | |
| Beardless Wheatgrass | | | | | | | | | | | |
| late veg | 33.1 | 10.3 | 12.9 | 72.7 | 18.8 | 3.4 | 6.2 | 41.4 | .41 | .17 | .20 |
| 4-26-80 | | | | | | | | | | | |
| mid bloom | 50.7 | 9.1 | 10.3 | 71.7 | 16.9 | 5.1 | 8.9 | 37.9 | .49 | .15 | .15 |
| 7-7-80 | | | | | | | | | | | |
| late bloom | 45.8 | 10.9 | 10.4 | 74.4 | 20.0 | 5.6 | 9.9 | 36.0 | .45 | .14 | .17 |
| 7-9-80 | | | | | | | | | | | |
| dough | 54.0 | 8.4 | 7.4 | 73.4 | 19.4 | 4.5 | 10.6 | 37.3 | .36 | .11 | .13 |
| 7-17-80 | | | | | | | | | | | |

SHERMAN**Big Bluegrass**

| | | | | | | | | | | | |
|-----------------------|------|------|------|------|------|-----|-----|------|------|-----|-----|
| late veg 5-26-80 | 34.5 | 7.9 | 11.8 | 74.0 | 20.1 | 3.1 | 7.6 | 40.2 | .42 | .18 | .20 |
| mid bloom 6-9-80 | 32.0 | 10.7 | 9.8 | 75.0 | 17.5 | 4.5 | 8.7 | 41.3 | .49 | .20 | .22 |
| late bloom 6-23-80 | 45.9 | 11.0 | 9.1 | 75.6 | 19.7 | 5.5 | 9.1 | 39.3 | .46 | .17 | .20 |
| dough 7-7-80 | 59.3 | 9.6 | 9.6 | 76.8 | 19.9 | 4.5 | 9.7 | 39.7 | 1.43 | .13 | .24 |

ALKAR**Tall Wheatgrass**

| | | | | | | | | | | | |
|----------------------|------|-----|-----|------|------|-----|-----|------|-----|-----|-----|
| late bloom 8-3-80 | 48.1 | 7.8 | 6.4 | 78.1 | 19.0 | 4.0 | 8.8 | 43.3 | .40 | .18 | .11 |
|----------------------|------|-----|-----|------|------|-----|-----|------|-----|-----|-----|

CASCADE**Birdsfoot Trefoil**

| | | | | | | | | | | | |
|----------------------|------|------|------|------|-----|-----|------|------|------|-----|-----|
| mid bloom 6-20-80 | 16.7 | 15.7 | 18.8 | 48.7 | 7.2 | 2.0 | 12.4 | 24.5 | 1.69 | .80 | .26 |
|----------------------|------|------|------|------|-----|-----|------|------|------|-----|-----|

ALTA***Tall Fescue**

| | | | | | | | | | | | |
|------------------------|-----|-----|------|------|------|-----|-----|------|-----|-----|-----|
| early bloom 5-14-73 | --- | --- | 11.8 | --- | 18.4 | 0.9 | 5.6 | 35.8 | -- | -- | -- |
| full bloom 6-14-80 | --- | 6.9 | 7.2 | 64.0 | 15.2 | 1.1 | 4.6 | 40.2 | .81 | .35 | .11 |
| 8-28-73 + 10-8-73 | --- | --- | 16.3 | 51.4 | 15.6 | 1.9 | 5.3 | 25.9 | -- | -- | -- |

*** Not a Pullman PMC release**

1 **Ash** inversely correlates with forage quality

2 **Crude protein** correlates with forage quality

3 **NDF** (Neutral Detergent Fiber) inversely correlates with forage quality

NDF = cellulose + hemicellulose + lignin + ash +/- error

4 **Hemicellulose** inversely correlates with forage quality

Hemicellulose partially digestible depending on type livestock, lignin content, etc.

5 **Acid Insoluble Ash** inversely correlates with forage quality

6 **Lignin** inversely correlates with forage quality

7 **Cellulose** inversely correlates with forage quality

Table 3. Forage quality of 'Nordan' and 'P-27' sampled at three stages of growth in 1992.
CULTIVAR

| STAGE OF GROWTH DATE | CRUDE PROTEIN | ACID DETERGENT FIBER (ADF) | ACID DETERGENT LIGNIN (ADL) |
|-----------------------------|------------------|-------------------------------|--------------------------------|
| ------(percent)----- | | | |
| NORDAN | | | |
| Crested Wheatgrass | | | |
| early vegetative 3-24-92 | 24.2 | 26.9 | 6.0 |
| late vegetative 5-1-92 | 13.2 | 32.6 | 4.6 |
| seed filling 6-8-92 | 6.5 | 37.0 | 12.4 |
| P-27 | | | |
| Siberian Wheatgrass | | | |
| early vegetative 3-24-92 | 18.2 | 32.2 | 7.6 |
| late vegetative 5-1-92 | 16.9 | 29.4 | 4.9 |
| seed filling 6-8-92 | 10.9 | 36.4 | 9.5 |